

**Jeffrey B. Witty, M.D.**  
Orthopaedic Surgery and Sports Medicine  
Acadiana Orthopaedic Center at Lafayette General  
4212 W. Congress Street  
Suite 1600  
Lafayette, LA 70506  
Phone: (337) 216 - 9018  
Fax: (337) 216 - 9143



**Isolated Grade II MCL Protocol: In brace unless removed by physician.**

First 48 – 72 hrs

- Rest, cryotherapy/ice, compression (tubigrip), elevation
- Hinged knee brace locked in extension if very painful / swollen and guarding – not usually for longer than a few days.
- PO NSAID

Brace:

- Duration of use based on clinical exam and progression with protocol.
- Use brace initially day and night (except bathing) until cleared for removal by MD.

After Resolution of Swelling:

- Begin gentle range of motion.
- E-stim, Hivamat, quad sets, straight leg raises
- Partial weight bearing once swelling/pain controlled and quickly advance to full weight bearing as pain allows
- Start normal walking once gait normalized/no limp and no pain and knee ROM at least 90 degrees
- Start progressive gentle closed chain exercise if pain allows.

After Gait Normalized and ROM at Least 90 Degrees:

- Continue progressive resistance closed kinetic chain exercises as comfort allows
- Stationary bike

Note: Will see physician prior to clearance for running / or drills OR please contact physician to discuss progress prior to beginning running or drills.

- Once WBAT, progress light jogging, progress agility (see below), static proprioceptive, neuromuscular, and sport specific drills

Once able to jog/run at 75% max speed and pain-free:

- Begin figure 8 drills at 20 yards, then progress to 10 yard figure 8 drills
- If pain-free, progress to cutting drills at 45 degrees, then 90 degree cutting drills
- Goal to achieve full ROM and approximately 80% strength of opposite side before returning to sport participation
- Continue to strengthen and dynamic proprioception until patient has achieved confidence to return to play

**Jeffrey B. Witty, M.D.**  
Orthopaedic Surgery and Sports Medicine  
Acadiana Orthopaedic Center at Lafayette General  
4212 W. Congress Street  
Suite 1600  
Lafayette, LA 70506  
Phone: (337) 216 - 9018  
Fax: (337) 216 - 9143



- Bracing up to 6 weeks after injury and a functional brace may be used during contact or high-risk activity.

Clear for Attempt to Return to Full Participation After:

- Normal ligamentous exam
- No pain with sport/activity agilities

Functional Running Program:

\*Begin after full range of motion, full weightbearing without pain, and pain-free jogging has been achieved

- Jog 1 mile
- Five successive 80 yard sprints at ½ speed
- Five successive 80 yard sprints at ¾ speed
- Five successive 80 yard sprints at full speed
- Five zigzag sprints at ½ speed
- Five zigzag sprints at full speed

Note: Return to play may take up to 3 – 8 weeks

Adapted from:

Noyes' Knee Disorders

Giannotti BF. American Journal of Sports Medicine 2006.

Edson CJ. Sport Med Arthrosc Rev 2006.

Reider B. American Journal of Sports Medicine 1994.